

Polarization of Comet Arend-Roland

WHEN, last May, Comet Arend-Roland became a bright naked-eye object in the northern hemisphere, we carried out a series of photoelectric measures of the polarization of its coma at different phase angles. The technique employed utilized a piece of 'Polaroid' attached to a rotatable six-position disk contained in the photometer. This enables measurements of intensity to be made as the 'Polaroid' is turned through 60 deg. for each position of the disk. Standard *B* and *V* filters employed in the *UBV* system, together with interference filters having peak transmissions centred on 5890 Å., 5000 Å., 4700 Å. and 4300 Å. and half-widths of 80 Å. were used for isolating the different wave-lengths. Each of the three circular apertures behind the 'Polaroid' transmitted 3.7 square minutes of arc of the sky.

We have calculated the amount of polarization and plane of polarization, from at least six sets of measures, each day, for radiation transmitted by each filter, a set of deflexions being considered as those obtained through three successive positions of the 'Polaroid'. Since, in each position of the 'Polaroid', the light reaches the photomultiplier through a diaphragm and area of the 'Polaroid' different from that of the previous position, we determined correction factors by measures of bright nearby stars, and applied these to the cometary measures. The orientation of the 'Polaroid' was determined by a laboratory arrangement, and we believe that the error in transferring this plane of reference to the telescope does not exceed 1 deg. The probable error of the value of percentage polarization is estimated to be less than ± 1 per cent and that of a value of the plane of polarization to be ± 5 deg. The values derived for 5890 Å., 5000 Å., and 4700 Å. are given in Table 1. These values cover a limited range of phase angle because the comet, after May 8, was beyond the range of accurate measurement with our interference filter - 'Polaroid' combination.

The measures of polarization through the *B* and *V* filters cover a larger phase angle than with the interference filters. Table 2 contains values of the amount of polarization and the plane of polarization for each day obtained through the *V*-filter, along with those of phase angle and position angle of the intensity equator that were calculated from the ephemeris of Candy¹. A general decrease in

Table 1

Date May 1957	Percentage polarization and position angle of plane of polarization		
	5890 A.	5000 A.	4700 A.
4-658	17.5 103°	—	17.9 107°
5-641	17.4 116°	18.0 112°	20.4 114°
6-648	19.7 119°	15.0 117°	17.3 119°
7-655	12.9 121°	14.1 124°	16.8 122°
8-660	20.5 123°	20.0 123°	14.5 —

amount of polarization with phase angle, along with superimposed erratic fluctuations from day to day of the order of 2-3 per cent, is seen. Assuming that the relation between amount of polarization and phase angle is given by

$$P_0 = \frac{P_{90} \sin^2 \theta}{1 + P_{90} \cos^2 \theta}$$

where P_0 and P_{90} are the values at phase angles 0° and 90° respectively, we have calculated the values of P_{90} for the comet through the different filters. These are seen to be consistent among themselves.

The 5890-A., 5000-A., 4700-A. interference filters were chosen to isolate the Swan bands present in the spectrum of the comet. A check on the continuum was maintained with the aid of the 4300-A. filter. The V -filter transmits, in the yellow and green regions, the C_2 radiations and possible sodium D -line emission together with the continuum of reflected sunlight. Hence, a value of the polarization of the coma determined through the V -filter represents a weighted mean of the percentage of polarization of the emission bands as well as of the continuum. The equality in

Table 2

Date May 1957	Percentage polariza- tion	Position angle of plane of polarization	Phase angle	Position angle of intensity- equator
4-658	18.9	104°	76° 24'	37° 48'
5-641	18.3	111°	73° 47'	40° 01'
6-648	19.9	116°	71° 12'	42° 12'
7-655	17.0	119°	68° 48'	44° 16'
8-660	14.5	120°	66° 36'	46° 13'
11-656	12.8	129°	60° 41'	51° 21'
14-662	14.0	148°	55° 45'	55° 25'
15-662	15.3	135°	54° 16'	56° 32'
22-647	9.6	147°	45° 44'	62° 00'
26-650	6.2	148°	41° 54'	63° 37'

the values of P_{90} derived through the different filters used, indicates the dominant part played by the continuum. We have had no access to spectra of the Comet taken on these days, but judging from Wellman's² description of the spectrum as it appeared on April 23, the emission may have been very weak during the period we observed. The polarization of Comet Arend-Roland is thus similar to that of Comet Paraskevopoulos (1941c) studied by Ohman³.

Within the errors of measurement, the planes of polarization at any one time of observation obtained through the different filters are all identical. Table 2 shows that the change in plane of polarization with time is linear, though it is more rapid than the change in position angle of the intensity-equator during the same time interval.

M. K. VAINU BAPPU
S. D. SINVAHAL

Uttar Pradesh State Observatory,
Naini Tal.
Aug. 6, 1957.

- ¹ Candy, M. P., Int. Astron. Union Circ. No. 1585 (Feb. 20, 1957).
² Wellman, P., Int. Astron. Union Circ. No. 1596 (April 29, 1957).
³ Ohman, Y., *Stockholm Annals*, 13, No. 11 (1941).