

# Interstellar Extinction in 20 Open Star Clusters

Geeta Rangwal<sup>1,4</sup>, R. K. S. Yadav<sup>2</sup>, Alok K. Durgapal<sup>1</sup> and D. Bisht<sup>3</sup>

<sup>1</sup>Center of Advanced Study, Department of Physics, D. S. B. Campus, Kumaun University Nainital 263002, India

<sup>2</sup>Aryabhata Research Institute of Observational Sciences, Manora Peak, Nainital 263002, India

<sup>3</sup>Astronomy and Astrophysics Division, Physical Research Laboratory, Ahmedabad 380009, Gujarat, India

<sup>4</sup>Email: [geetarangwal91@gmail.com](mailto:geetarangwal91@gmail.com)

(RECEIVED August 16, 2017; ACCEPTED November 21, 2017)

## Abstract

The interstellar extinction law in 20 open star clusters namely, Berkeley 7, Collinder 69, Hogg 10, NGC 2362, Czernik 43, NGC 6530, NGC 6871, Bochum 10, Haffner 18, IC 4996, NGC 2384, NGC 6193, NGC 6618, NGC 7160, Collinder 232, Haffner 19, NGC 2401, NGC 6231, NGC 6823, and NGC 7380 have been studied in the optical and near-IR wavelength ranges. The difference between maximum and minimum values of  $E(B - V)$  indicates the presence of non-uniform extinction in all the clusters except Collinder 69, NGC 2362, and NGC 2384. The colour excess ratios are consistent with a normal extinction law for the clusters NGC 6823, Haffner 18, Haffner 19, NGC 7160, NGC 6193, NGC 2401, NGC 2384, NGC 6871, NGC 7380, Berkeley 7, Collinder 69, and IC 4996. We have found that the differential colour-excess  $\Delta E(B - V)$ , which may be due to the occurrence of dust and gas inside the clusters, decreases with the age of the clusters. A spatial variation of colour excess is found in NGC 6193 in the sense that it decreases from east to west in the cluster region. For the clusters Berkeley 7, NGC 7380, and NGC 6871, a dependence of colour excess  $E(B - V)$  with spectral class and luminosity is observed. Eight stars in Collinder 232, four stars in NGC 6530, and one star in NGC 6231 have excess flux in near-IR. This indicates that these stars may have circumstellar material around them.

Keywords: dust, extinction – ISM: general – open clusters and associations: general

## 1 INTRODUCTION

Interstellar dust is an important component of the interstellar medium. It is the remnant of star formation and stellar evolution processes. Interstellar dust grains can transmit, redirect, and transmute the starlight (Fitzpatrick 2004). Due to this property of dust grain, the actual distances and magnitude determination of astronomical objects is often very difficult (Pandey et al. 2002). For this reason, it is very important to have knowledge of interstellar dust in the line of sight of the objects under study. The interstellar extinction is caused by two distinct sources, a diffuse component associated by the general interstellar medium and a variable component associated with localised region of higher mean density (Martin & Whittet 1990; Turner 1994). Studies of the variable component gives information about the composition of stars. Extinction properties of dust are wavelength dependent, so investigation of interstellar extinction law is the best way to determine the dust properties (Yadav & Sagar 2001; Fitzpatrick 2004). Young open star clusters are the ideal objects for this kind of study because they may contain gas and dust around early type (*O*, *B*, and *A*) stars (Hayashi 1970; Lar-

son 1973; McNamara 1976, Warner, Strom, & Strom 1979; Yadav & Sagar 2001).

Trumpler studied the interstellar dust for the first time in 1920 and then it was studied by many investigators. Krelowski and Strobel (1983) analysed interstellar extinction law in two stellar aggregates namely, Sco OB2 and Per OB1 and found different extinction curves for these two aggregates and their differences are large in far UV wave band. They concluded that the differences in FUV (Far-Ultraviolet) fluxes for these fields is due to the circumstellar dust and also confirmed the presence of two populations of grains. Kizskurno et al. (1984) investigated 20 large OB associations at different galactic positions and found similar colour excess curves in the FUV range. They also concluded that these complexes are obscured by same type of interstellar material. Sagar (1987) studied interstellar extinction in 15 open star clusters and found non-uniform extinction across 10 of the clusters. When the value of the inferred colour excess varies from one part of cluster region to another one, this is known as non-uniform extinction. The possible reasons behind the non-uniform extinction has been discussed by many investigators. Sagar (1987) and Sagar & Qian (1989) stated that this

non-uniformity is due to the hot and ionised circumstellar dust around the stars. Samson (1975), McCuskey & Hauk (1964), and Stone (1977) proposed that dust shells around stars and local dust clouds lying in the cluster direction is responsible for non-uniform extinction. Tapia et al. (1988) studied the extinction law in clusters using 200 stars located within four open clusters situated at same distance from the Sun. They found that these clusters have different reddening values and that they follow the normal extinction law for  $\lambda > 0.55 \mu\text{m}$ . They also interpreted the anomalies of the extinction law in then  $U$  and  $B$  bands because of the presence of dense intra-cluster dust cloud of different grain properties in comparison with normal dust grains. Cardelli, Clayton, & Mathis (1989) derived the mean extinction law for the wavelength range  $0.12 \mu\text{m} \leq \lambda \leq 3.5 \mu\text{m}$ . This extinction law is applicable for the diffuse as well as the dense interstellar medium and depends only on one parameter  $R_V$ , the total to selective extinction ratio.

Pandey, Mahra, & Sagar (1990) found that differential extinction which is due to a non-uniform distribution of star-forming material decreases systematically with the age of the cluster. Yadav & Sagar (2001) studied interstellar extinction law in 15 open star clusters and found the presence of non-uniform extinction in all the clusters. They also found that most of the clusters follow the normal extinction law for  $\lambda < \lambda_J$ , more specifically that the value of the colour excess for these clusters is close to the standard value derived for the general interstellar medium in this waveband, but show anomalous behaviour for  $\lambda \geq \lambda_J$ . Pandey et al. (2002) studied 14 open clusters located around  $l \sim 130^\circ$  and found that dust along these objects follow the normal interstellar extinction law for  $\lambda \geq \lambda_J$ , but for shorter wavelengths, it depends upon  $R_{\text{cluster}}$  (total-to-selective absorption in the cluster region). Joshi (2005) studied the distribution of interstellar matter near the Galactic plane on the basis of 722 open star clusters and found that  $\sim 90\%$  of the absorbing material lies within  $-5^\circ \leq b \leq 5^\circ$  of the Galactic plane. Nishiyama et al. (2005) studied the interstellar extinction law in  $J$ ,  $H$ , and  $K_S$  bands towards the Galactic centre in the region  $l \leq 2^\circ.0$  and  $0^\circ.5 \leq b \leq 1^\circ.0$  using Red Clump (RC) stars and concluded that the extinction law is not universal even in the infra-red band.

In the light of above discussions, we can say that there are numerous studies available regarding the interstellar extinction in the direction of different objects. But, yet there are number of clusters for which less information is available about the distribution of dust and gas in our Galaxy. In order to study the distribution of dust and gas in the Milky Way, we need to consider different young open star clusters. Therefore, we have in this study considered 20 open star clusters to study interstellar extinction law in the optical and near-IR bands.

The structure of the article is as follows. Section 2 describes the data sources and cluster selection criteria. Section 3 describes the data analysis while Section 4 represents the conclusions of the present study.

## 2 CLUSTER SELECTION AND DATA SET

To study the extinction properties in the clusters, we have selected 20 young open star clusters from the WEBDA database, which have age younger than 50 Myr. The selection criteria of these clusters is based on the availability of  $UBVRI$  photometric data and previous studies on extinction. These clusters have been considered first time for interstellar extinction study. These clusters have distances ranging from 0.44 to 6.3 kpc. In young open star clusters, the early type stars may have dust and gas around them. Therefore, for the present analysis, we have selected only early type stars ( $O$ ,  $B$ , and  $A$ ) in each cluster. These stars have been selected using colour–colour and colour–magnitude diagrams (CMDs). The CMDs of the clusters are shown in Figure 1. The crosses represent the selected stars for the present analysis, while gray points are the remaining stars of the clusters. The  $UBVRI$  data for these clusters are taken from the WEBDA (<http://www.univie.ac.at/webda/>) database, while  $JHK$  data are taken from 2MASS (Two Micron All Sky Survey) catalogue. These clusters are distributed along the galactic plane with longitude from  $6^\circ.08$  to  $336^\circ.70$ . Almost all the clusters under study have  $UBVRIJHK$  data. The optical data set have error  $\sim 0.01$  mag in  $V$  and  $\sim 0.02$  mag in  $(B - V)$ ,  $(V - R)$ , and  $(V - I)$  while  $\sim 0.03$  mag in  $(U - B)$ . The  $JHK$  data set have error  $\sim 0.05$  mag in  $J$  mag. The spectroscopic data is taken from the WEBDA database wherever available. Due to the unavailability of spectroscopic data for many stars, photometric Q-method (Johnson & Morgan 1953) have been used. The maximum difference between the spectral type as identified by Q-method and that of available spectral type data is about two spectral sub-classes. The general information about the selected open star clusters is listed in Table 1.

## 3 ANALYSIS OF OBSERVATIONAL DATA

For the study of interstellar extinction law, we have derived the  $E(U - V)$ ,  $E(B - V)$ ,  $E(V - R)$ ,  $E(V - I)$ ,  $E(V - J)$ ,  $E(V - H)$ , and  $E(V - K)$  colour excesses for early type stars. We have calculated these excesses by comparing the observed colours with their intrinsic ones. The intrinsic colours are derived from the MKK spectral type luminosity class colour relation given by FitzGerald (1970) for  $(U - V)$  and  $(B - V)$ , by Johnson (1966) for  $(V - R)$  and  $(V - I)$  and by Koornneef (1983) for  $(V - J)$ ,  $(V - H)$ , and  $(V - K)$ . The number of stars included in the study for a particular cluster are listed in the Table 1.

### 3.1. Non-uniform extinction

To study the presence of non-uniform extinction in each cluster, we have calculated  $\Delta E(B - V) = E(B - V)_{\text{max}} - E(B - V)_{\text{min}}$ , where  $E(B - V)_{\text{max}}$  and  $E(B - V)_{\text{min}}$  are the average  $E(B - V)$  of four highest and four lowest values of  $E(B - V)$ . The values of  $\Delta E(B - V)$  along with the central coordinates of each clusters are listed in the Table 2. The

**Table 1.** General information about the clusters under study taken from WEBDA website.  $N_S$  denotes the number of stars used in the analysis.


IAU number	Clusters	$l$ (°)	$b$ (°)	Distance (kpc)	log(age) (yr)	$N_S$
C0150+621	Berkeley 7	130.13	0.37	2.57	6.60	40
C0532+099	Collinder 69	195.05	-12.00	0.44	7.05	08
C1108-601	Hogg 10	290.80	0.07	1.77	6.78	26
C0716-248	NGC 2362	238.18	-5.54	1.39	6.91	17
C2323+610	Czernik 43	112.84	0.13	2.50	7.70	24
C1801-243	NGC 6530	6.08	-1.33	1.33	6.87	33
C2004+356	NGC 6871	72.64	2.08	1.57	7.00	83
C1040-588	Bochum 10	287.03	-0.32	2.03	6.85	11
C0750-262	Haffner 18	243.11	0.44	6.03	6.00	56
C2014+374	IC 4996	75.36	1.31	1.73	6.95	21
C0722-209	NGC 2384	235.39	-2.41	2.12	6.00	10
C1637-486	NGC 6193	336.70	-1.57	1.15	6.77	20
C1817-162	NGC 6618	15.09	-0.74	1.30	6.00	11
C2152+623	NGC 7160	104.01	6.45	0.79	7.28	14
C1042-59	Collinder 232	287.49	-0.54	2.99	6.67	26
C0750-261	Haffner 19	243.08	0.52	5.09	6.93	41
C0727-138	NGC 2401	229.66	1.85	6.30	7.40	33
C1650-417	NGC 6231	243.46	1.18	1.24	6.84	123
C1941+231	NGC 6823	59.40	-0.14	1.89	6.82	77
C2245+578	NGC 7380	107.14	-0.88	2.22	7.08	88

**Table 2.** The values of mean  $E(B - V)$ ,  $E(B - V)_{\min}$ ,  $E(B - V)_{\max}$ , and  $\Delta E(B - V)$  of each cluster are listed with their central coordinates.

Cluster	Central coordinates		$\overline{E(B - V)}$ (Mag)	$E(B - V)_{\min}$ (Mag)	$E(B - V)_{\max}$ (Mag)	$\Delta E(B - V)$ (Mag)
	$\alpha_{2000}$	$\delta_{2000}$				
Berkeley 7	01:54:12	+62:22:00	0.81	0.66	1.02	0.36
Collinder 69	05:35:06	+09:56:00	0.15	0.10	0.21	0.11
Hogg 10	11:10:42	-60:24:00	0.40	0.30	0.49	0.19
NGC 2362	07:18:41	-24:57:18	0.11	0.08	0.17	0.09
Czernik 43	23:25:48	+61:19:11	0.61	0.52	0.73	0.21
NGC 6530	18:04:31	-24:21:30	0.34	0.27	0.40	0.13
NGC 6871	20:05:59	+35:46:36	0.44	0.32	0.56	0.24
Bochum 10	10:42:12	-59:08:00	0.41	0.34	0.48	0.14
Haffner 18	07:52:39	-26:23:00	0.40	0.24	0.53	0.29
IC 4996	20:16:30	+37:38:00	0.67	0.56	0.77	0.21
NGC 2384	07:25:10	-21:01:18	0.27	0.22	0.31	0.09
NGC 6193	16:41:20	-48:45:48	0.47	0.39	0.53	0.14
NGC 6618	18:20:47	-16:10:18	0.39	0.31	0.48	0.17
NGC 7160	21:53:40	+62:36:12	0.39	0.32	0.48	0.16
Collinder 232	10:44:59	-59:33:00	0.41	0.30	0.52	0.22
Haffner 19	07:52:47	-26:17:00	0.38	0.25	0.48	0.23
NGC 2401	07:29:24	-13:58:00	0.35	0.29	0.41	0.12
NGC 6231	16:54:10	-41:49:30	0.46	0.35	0.63	0.28
NGC 6823	19:43:09	+23:18:00	0.86	0.66	1.04	0.38
NGC 7380	22:47:21	+58:07:54	0.56	0.40	0.69	0.29


factors which can produce dispersion in  $\Delta E(B - V)$  other than non-uniform extinction can cause maximum dispersion  $\sim 0.11$  mag (Burki 1975; Sagar 1985; Sagar 1987; Pandey et al. 1990; Yadav & Sagar 2001). Therefore, the value of  $\Delta E(B - V)$  greater than 0.11 mag can be considered as non-uniform extinction in the cluster. In the present study, except

three clusters namely, NGC 2362, NGC 2384, and Collinder 69, all are having  $\Delta E(B - V)$  value more than 0.11 mag. So this analysis confirms the presence of non-uniform extinction in all clusters except NGC 2362, NGC 2384, and Collinder 69. In case of NGC 6530, we found a result similar as that reported by Sagar (1987).




**Figure 1.** The  $V$ ,  $(B - V)$  colour-magnitude diagrams for all 20 clusters under study. The gray points represent all stars of the cluster and black crosses represent the stars used for the present study.

In addition to this, we have also analysed the extent of non-uniform extinction by plotting the histograms of  $E(B - V)$  as shown in Figure 2 for each cluster. Histograms show that all the clusters have a wide range in  $E(B - V)$  values. This indicates that all the clusters have different amounts of non-



**Figure 2.** The dispersion of  $E(B - V)$  shows the presence of non-uniform extinction for the clusters under study.


uniform extinction and dust is non-uniformly distributed over the cluster region. In the following sections, we explore the possible reasons for the presence of this type of distribution in  $E(B - V)$ .



**Figure 3.** The colour excesses  $E(U - V)$ ,  $E(V - R)$ ,  $E(V - I)$ ,  $E(V - J)$ ,  $E(V - H)$ ,  $E(V - K)$  are plotted against  $E(B - V)$  for the clusters Berkeley 7, NGC 6823, Collinder 232, and NGC 6530.

### 3.2. Extinction law

To investigate the interstellar extinction law towards the clusters, we have plotted the colour excesses  $E(U - V)$ ,  $E(V - R)$ ,  $E(V - I)$ ,  $E(V - J)$ ,  $E(V - H)$ , and  $E(V - K)$  against  $E(B - V)$  in Figures 3–7. The solid line in the plots is the least-square fit to the data points. The slope of the colour excess ratios along with their error are listed in Table 3. The theoret-



**Figure 4.** Same as Figure 3 for the clusters Haffner 19, NGC 7160, NGC 6193, and Bochum 10.

ical values of the colour excess ratios are taken from Pandey et al. (2002). By comparing the observed colour excess ratios with the theoretical one, following conclusions can be drawn:

- (i) Twelve clusters namely NGC 6823, Haffner 18, Haffner 19, NGC 7160, NGC 6193, NGC 2401, NGC 2384, NGC 6871, NGC 7380, Berkeley 7, Collinder



**Table 3.** The derived and normal colour excess ratios for all the clusters under study.

Cluster	$\frac{E(U - V)}{E(B - V)}$	$\frac{E(V - R)}{E(B - V)}$	$\frac{E(V - I)}{E(B - V)}$	$\frac{E(V - J)}{E(B - V)}$	$\frac{E(V - H)}{E(B - V)}$	$\frac{E(V - K)}{E(B - V)}$
Normal value	1.72	0.60	1.25	2.30	2.58	2.78
Berkeley 7	1.72±0.03	0.67±0.02	1.18±0.05	1.98±0.13	2.48±0.13	2.61±0.13
NGC 6823	1.74±0.03	0.67±0.05	1.39±0.16	2.10±0.23	2.44±0.18	2.64±0.22
Collinder 232	1.88±0.06	0.62±0.05	1.37±0.13	2.55±0.37	2.98±0.52	3.18±0.57
NGC 6530	1.87±0.17	0.81±0.15	1.57±0.16	3.21±0.13	3.55±0.46	4.19±0.48
Haffner 19	1.73±0.09	0.61±0.04	1.42±0.10	2.10±0.25	2.91±0.17	3.10±0.28
NGC 7160	2.00±0.19	0.64±0.20	1.35±0.23	2.14±0.29	2.09±0.35	2.50±0.27
NGC 6193	1.86±0.18	0.68±0.07	1.59±0.18	2.03±0.37	2.68±0.44	2.86±0.54
Bochum 10	2.38±0.30	0.64±0.15	1.40±0.55	2.17±0.64	2.37±0.60	3.20±0.69
Haffner 18	1.76±0.03	0.61±0.03	1.30±0.09	2.06±0.23	2.78±0.13	2.93±0.23
NGC 2362	2.25±0.23	0.47±0.13	1.15±0.28	2.41±0.86	2.68±0.78	3.94±0.74
Hogg 10	1.99±0.06	0.60±0.08	1.39±0.09	2.07±0.25	2.27±0.38	2.66±0.45
NGC 2401	1.84±0.14	0.77±0.11	1.53±0.18	2.62±0.36	2.88±0.28	2.63±0.32
IC 4996	1.75±0.10	0.33±0.06	1.86±0.12	1.70±0.32	2.18±0.42	2.64±0.45
NGC 2384	1.65±0.21	0.33±0.28	2.34±0.87	2.33±0.58	2.29±0.39	2.76±0.33
Czernik 43	1.78±0.05	0.54±0.10	*	1.55±0.23	2.10±0.24	2.22±0.26
NGC 6871	1.68±0.05	*	*	2.02±0.27	2.37±0.19	2.50±0.20
NGC 6618	1.91±0.06	*	*	1.93±0.21	2.40±0.52	2.83±0.82
NGC 7380	1.76±0.03	*	*	2.11±0.16	2.68±0.15	2.78±0.19
NGC 6231	1.85±0.04	*	1.24±0.03	2.15±0.16	2.64±0.14	2.75±0.16
Collinder 69	1.79±0.10	*	*	2.45±0.57	2.30±0.36	3.86±0.36

69, and IC 4996 follow the normal extinction law in optical and near-IR region within  $2\sigma$ , where  $\sigma$  is the standard deviation of data points. This shows the expected behaviour of the normal interstellar law in the direction of these clusters and hence presumably dust with normal characteristics in shape, size, and composition (Tapia et al. 1991). Similar results have also been discussed in Yadav & Sagar (2001) for the clusters NGC 1893, NGC 2244, Tr 37, and Be 68.

- (ii) Six clusters namely Bochum 10, NGC 2362, Collinder 232, Hogg 10, NGC 6618, and NGC 6231 show anomalous behaviour in  $\lambda < \lambda_J$ . These clusters have colour excess ratios greater than that of the normal value in optical band. This indicates that extinction law is different than for the normal extinction law. This also implies that dust particles in the direction of these clusters have smaller cross section than the normal ones (Turner 1994).
- (iii) Two clusters NGC 6530 and Czernik 43 show anomalous behaviour for  $\lambda \geq \lambda_J$ . NGC 6530 have large colour excess ratios, while Czernik 43 have less ratios than the normal one.

The extinction law around early type stars can be understood by the model given by Seab & Shull's (1983) for dust grain processing due to passage of shock waves. Results of this model are naturally dependent on the physical conditions of the intracluster material (Sagar & Qian 1993). Small inhomogeneities in them would therefore result in large differences in the observed parameters. Consequently, depending upon the physical conditions of intracluster material, grain size

distribution resulting from the interactions of strong radiations of hot O and B stars can be either normal or shifted by a small amount in either direction compared to that of normal sized particles. The anomaly depends on the direction of the shift, which ultimately depends on the grain destruction mechanism.

### 3.3. Variation of colour excess with age of the clusters

It is believed that stars are formed in the dense clouds of dust and gas, by contracting under their own gravitational attraction (Yorke & Krugel 1977; Pandey et al. 1990). Not all of this gas or dust becomes part of star but also used in the formation of planets, asteroids, comets, etc. or remains as dust. So this remnant dust is either used in the star formation or blown away by the high radiation pressure of embedded or nearby stars (Pandey et al. 1990). Therefore, young open star clusters should have some amount of dust and gas which gradually decreases with time. It means there should be a relationship between age of the star cluster and the interstellar extinction. Pandey et al. (1990) studied the variation of non-uniform extinction  $\Delta E(B - V)$  with age for 64 open star clusters and found that extinction decreases with the age of the clusters and calculated the gas removal time larger than  $10^8$  yrs. As  $\Delta E(B - V)$  represents the measure of non-uniform extinction, so we have plotted the age of each cluster with  $\Delta E(B - V)$  to see the relationship between age of cluster and the presence of non-uniform extinction. This plot is shown in Figure 8. The values of  $\Delta E(B - V)$  of the clusters under study are listed in the Table 2 and the ages of the clusters are taken from Table 1. For this study, we have also included




Figure 5. Same as Figure 3 for the clusters Haffner 18, NGC 2362, Hogg 10, and NGC 2401.

the clusters studied by Yadav & Sagar (2001). To see the variation of mean  $\Delta E(B - V)$  with age, we have binned the data into several groups with an age interval of 0.5 Myr. After binning the data, we have calculated the mean of  $\Delta E(B - V)$  and its standard deviation, which are plotted in Figure 8. The variation of  $\Delta E(B - V)$  with age indicates that a large amount of obscuring material is present in younger age clusters. For older clusters, the mean value of  $\Delta E(B - V)$  is small. So

PASA, 34, e068 (2017)  
doi:10.1017/pasa.2017.64





Figure 6. Same as Figure 3 for the clusters IC 4996, NGC 2384 Czernik 43, and NGC 6231.

we can conclude that the extent of non-uniform extinction decreases with age. Dust and gas may be blown away from the older clusters due to several reasons as discussed in Yadav & Sagar (2001). A similar analysis was also done by Yadav & Sagar (2001) and found the similar relationship between  $\Delta E(B - V)$  and age of the clusters.

### 3.4. Spatial variation of colour excess


To study the spatial variation of colour excess  $E(B - V)$ , we have divided the whole region of the clusters into small



**Figure 7.** Same as [Figure 2](#) for the clusters NGC 6618, NGC 7380, Collinder 69, and NGC 6871.

square boxes. The clusters Collinder 69, Collinder 232, Czernik 43, NGC 6530, Bochum 10, NGC 2384, NGC 6618, and NGC 7160 were not considered for this study due to insufficient number of stars and smaller area. The area of the clusters Berkeley 7, Haffner 18, Haffner 19, Hogg 10, IC 4996, NGC 2401, NGC 6193, NGC 6231, and NGC 6823 is divided into small boxes of area  $5 \text{ arcmin} \times 5 \text{ arcmin}$ . For the clusters NGC 6871 and NGC 7380, the square boxes are of  $10 \text{ arcmin} \times 10 \text{ arcmin}$  due to the available larger area. We have calculated the mean value of  $E(B - V)$  and its standard deviation for the stars in each boxes. The close inspection of these values provides a clue that there is no significant variation of the mean colour excess values with the positions in the cluster area. A small variation of mean  $E(B - V)$  with position in NGC 6193 is observed and listed in [Table 4](#). In NGC 6193, the value of  $E(B - V)$  decreases from east to

PASA, 34, e068 (2017)  
doi:[10.1017/pasa.2017.64](https://doi.org/10.1017/pasa.2017.64)



**Figure 8.** Variation of  $\Delta E(B - V)$  with the cluster age. The open circles denote data points from present work and the filled circles denotes data points from [Yadav & Sagar \(2001\)](#). *Note:* Error bars denote the corresponding standard deviation of binned data.

west. A variation in  $E(B - V)$  with position has also been observed by [Sagar \(1987\)](#) for three clusters NGC 6611, IC 1805, and NGC 6530 and by [Yadav & Sagar \(2001\)](#) for Tr 14.

### 3.5. Spectral type variation of colour excess

To understand the relation between the interstellar extinction and the spectral type of the stars, we plotted  $E(B - V)$  against the spectral type of the stars with gray open points in [Figure 9](#). To see the significant variations, we have binned the stars in groups and calculated mean and standard deviation of  $E(B - V)$ . The plots of mean  $E(B - V)$  against spectral type are shown in [Figure 9](#) with black solid points. These plots clearly show that for most of the clusters there is no significant trend in  $E(B - V)$  with spectral class except Berkeley 7, Hogg 10, NGC 6871, and NGC 7380. For the clusters NGC 7380, NGC 6871, and Berkeley 7 initially  $E(B - V)$  remains constant, but after  $B6$  spectral class, it decreases. In Hogg 10,  $E(B - V)$  decreases continuously from  $O$  to  $A$  type stars.


### 3.6. Colour excess variation with luminosity

To study the variation of  $E(B - V)$  with the luminosity of stars, we have plotted the colour excess  $E(B - V)$  against the absolute magnitude of the stars. These plots are shown in [Figure 10](#) with gray open points. Apparent  $V$  magnitudes are converted into absolute magnitude  $M_V$  by the relation  $M_V = V - (5 \log d - 5 + A_V)$  where  $A_V = 3.25 E(B - V)$  and distances are taken from [Table 1](#). To see the clear variation in the colour excess, we have plotted the mean and standard deviation of  $E(B - V)$  against the absolute luminosity and shown in [Figure 10](#) with the black solid points with errors. We




**Table 4.** The spatial variation of  $E(B - V)$  across the open cluster NGC 6193. In each boxes, there is mean values of  $E(B - V)$  with their standard deviation for corresponding  $5 \times 5$  arcmin<sup>2</sup>. The numbers in the bracket represent the number of stars in that box.

$\Delta\alpha / \Delta\delta$	-5-0	0-5	5-10	10-15	15-20
-10--5	-	0.46(1)	-	-	-
-5-0	0.39±0.01(3)	-	0.49±0.04(5)	0.52±0.04(2)	-
0-5	0.40±0.01(2)	0.50(1)	0.48±0.01(5)	-	0.50(1)



**Figure 9.** Variation of  $E(B - V)$  with the spectral class of the stars. The gray open circles represent the data points while the black solid points are mean  $E(B - V)$ .



**Figure 10.** Variation of  $E(B - V)$  with the luminosity of the stars. The gray open circles represent the data points while the solid black points are mean  $E(B - V)$ .

grouped the stars to calculate the mean and standard deviation in colour excess. A close inspection of Figure 10 shows that there is no significant correlation of mean  $E(B - V)$  with luminosity for clusters Collinder 69, NGC 2384, NGC 6618,

**Table 5.** Near-IR flux excess in stars of cluster under study. Star numbers are taken from WEBDA.

Cluster	Stars	$V$ (Mag)	$E(V - J)$ (Mag)	$\Delta(V - H)$ (Mag)	$\Delta(V - K)$ (Mag)	SP type
Collinder 232	3	12.66	1.18	0.35	0.61	B3
	4	14.06	1.12	0.51	0.60	B9.5
	27	14.19	1.39	0.82	1.02	B9.5
	41	13.22	1.68	0.90	0.92	B8
	46	13.62	1.51	0.64	0.61	B8
	1 011	11.62	1.61	0.62	0.69	B3
	1 025	13.02	1.68	0.64	0.77	B3
	1 050	14.11	1.32	0.65	0.76	B9
NGC 6530	26	11.59	1.44	0.27	0.69	B5
	31	11.69	1.18	0.53	0.64	B6
	64	11.63	0.04	0.81	2.08	B6
	263	11.43	1.43	0.44	0.66	B5
NGC 6231	370	12.23	1.37	0.62	0.81	B8

NGC 7160, Haffner 18, Collinder 232, Czernik 43, NGC 2362, NGC 2401, Haffner 19, NGC 6823, NGC 2362, IC 4996, NGC 6231, and NGC 6618. However, in the clusters NGC 7380, NGC 6871, and Berkeley 7, a decreasing trend of  $E(B - V)$  with luminosity is observed after  $M_v \sim 0$  mag. In Hogg 10,  $E(B - V)$  decreases continuously with  $M_v$ .

### 3.7. Discussions of variable reddening

On the basis of the study presented in Sections 3.5 and 3.6, we divided the clusters sample into following three groups:

- (1) Clusters namely Berkeley 7, NGC 7380, NGC 6871 belong to group 1 show a dependence of  $E(B - V)$  on both spectral type and luminosity. Initially,  $E(B - V)$  remains constant up to B6 spectral class and  $M_v \sim 0$  mag luminosity but decreases as we go towards late type spectral class and fainter stars.
- (2) For the cluster Hogg 10 which is in group 2,  $E(B - V)$  continuously decreases with luminosity and temperature of the stars.
- (3) The clusters belonging to group 3 are Collinder 69, NGC 2384, NGC 6618, NGC 7160, Haffner 18, Collinder 232, Czernik 43, NGC 2362, NGC 2401, Haffner 19, NGC 6823, NGC 2362, IC 4996, NGC 6231, Hogg 10, and NGC 6618. For these clusters, no dependence of  $E(B - V)$  on either spectral type or luminosity is seen.



The observed dependence of  $E(B - V)$  on the spectral type and luminosity of the stars in groups 1 and 2 is due to the presence of circumstellar matter around them as discussed by Sagar (1987). In these groups, the value of  $E(B - V)$  is higher for early type and most luminous stars. Yorke & Krugel (1977) and Bhattacharjee & Williams (1980) discussed that relative mass of the relict envelope present around a newly formed massive star is positively correlated with the stellar

mass. Massive main sequence stars blow off their circumstellar material by emitting strong stellar winds and ultraviolet radiations. But the time required to blow off this circumstellar material is dependent on various parameters as discussed by Sagar (1987). Therefore, stars of groups 1 and 2 have their circumstellar material still present around them, whereas group 3 stars have blown their circumstellar envelope away.

### 3.8. Near-IR fluxes

The total interstellar extinction towards a star is the summation of three terms as stated by Yadav & Sagar (2001). These three terms are (i) interstellar extinction caused by the dust and gas present between the observer and the cluster, (ii) intra-cluster extinction caused by the dust and gas present inside the cluster field, and (iii) circumstellar extinction caused by the remnant discs or envelope around the young cluster members in which the disc is not completely dissipated.

Generally, it is assumed that the interstellar extinction caused by general interstellar medium between observer and the object follows the normal interstellar extinction law. So if there is an anomaly in the observed extinction law, then it may be due to the intra-cluster extinction or due to the circumstellar extinction or due to both. Since in the present study we have selected brighter stars in young clusters, so it may be possible that the observed anomaly in the interstellar extinction law may be due to the reasons listed in points (ii) and (iii) in above paragraph. To test these two possibilities, we have to use the colour excess  $E(V - J)$  instead of  $E(B - V)$  as stated by Smith (1987), Tapia et al. (1988), Sagar & Qian (1989, 1990), and Yadav & Sagar (2001) that the value of  $E(V - J)$  is independent of the properties such as chemical composition, shape, structure, and degree of alignment of the interstellar dust and cloud (Vashchinnikov & Il'in 1987; Cardelli, Clayton, & Mathis 1989). In order to study the reason behind the anomaly in the observed interstellar law, we have calculated the differences between the observed values of  $E(V - H)$  and  $E(V - K)$  based on spectral classification





**Figure 11.** Plots of near-IR flux excess in terms of  $\Delta(V-H)$  and  $\Delta(V-K)$  against the colour excess  $E(V-J)$ . The black line denotes the zero excess and the broken lines denote the extent of significant error.

and the derived colour excess values from  $E(V-J)$  assuming normal extinction law and has been plotted against the corresponding value of  $E(V-J)$  in Figures 11 and 12.

In the Figures 11 and 12,  $\Delta(V-H)$  and  $\Delta(V-K)$  are the differences of observed and derived colour excess, the continuous line denote the zero value of  $\Delta(V-H)$  and  $\Delta(V-K)$ . The errors in  $\Delta(V-H)$  and  $\Delta(V-K)$  mainly occur due to observational uncertainties in  $JHK$  magnitudes, inaccuracies in the calculation of  $E(V-J)$ , and errors in spectral classification. The maximum value of differences due to these factors can be  $\sim 0.5$  mag and denoted by dotted lines in Figure 11.

On close inspection of Figures 11 and 12, we can see that for most of the clusters, stars are concentrated around the zero value and inside the error line, which indicates that there is no near-IR excess fluxes in the stars of the clusters. In the clusters namely Collinder 232, NGC 6530, and NGC 6231



**Figure 12.** Plots of near-IR flux excess in terms of  $\Delta(V-H)$  and  $\Delta(V-K)$  against the colour excess  $E(V-J)$ . The black line denotes the zero excess and the broken lines denote the extent of significant error.

there are some stars which have larger value of  $\Delta(V-H)$  and  $\Delta(V-K)$  and they are lying outside the error boundary. These stars are listed in Table 5.

All the stars listed in Table 5 for the clusters Coll 232, NGC 6530 and NGC 6231 have larger values in  $\Delta(V-K)$ . It is possible that circumstellar material is present around these stars. We have detected near-IR fluxes for these stars for the first time. Strom, Strom, & Yost (1971) and Strom et al. (1972) noticed the presence of circumstellar material around NGC 2264-90 on the basis of near-IR fluxes. Smith (1987) and Yadav & Sagar (2001) have also found near-IR excess fluxes for the stars Tr 14-15, NGC 2264-165, and Tr 16-68.

## 4 CONCLUSIONS

In the present study, we analysed the interstellar extinction law in the direction of 20 open clusters younger than 50 Myr.

We have also studied the variation of colour excess  $E(B - V)$  with age, luminosity, spectral type, and position of stars in the cluster. The main results of this study are summarised as follows:

1. The histograms of  $E(B - V)$  show that gas and dust is non-uniformly distributed over the cluster area in each cluster. The value of  $\Delta E(B - V)$  confirms the presence of non-uniform extinction in all clusters under study except NGC 2362, NGC 2384, and Collinder 69. Our analysis shows that non-uniform extinction decreases with the age of the clusters. This indicates that dust and gas are still present in younger clusters while it has been blown away in older clusters by UV radiations of hot stars.
2. Derived colour excess ratios show that 12 clusters NGC 6823, Haffner 18, Haffner 19, NGC 7160, NGC 6193, NGC 2401, NGC 2384, NGC 6871, NGC 7380, Berkeley 7, Collinder 69, and IC 4996 have normal extinction laws in the optical and near-IR regions within  $2\sigma$  error. This indicates that dust and gas present in the clusters have normal size distributions. Six clusters Bochum 10, NGC 2362, Collinder 232, Hogg 10, NGC 6618, and NGC 6231 show anomalous behaviour in the optical band. Two clusters NGC 6530 and Czernik 43 show anomalous behaviour for  $\lambda \geq \lambda_j$ . This shows that the extinction law for these clusters is different from normal one.
3. Spatial variation of the colour excess  $E(B - V)$  has been found only for one cluster, NGC 6193, in the sense that  $E(B - V)$  is decreasing from east to west. For other clusters,  $E(B - V)$  is distributed non-uniformly over the cluster field.
4. A small variation in  $E(B - V)$  with spectral class and luminosity of stars have been found in the clusters Berkeley 7, NGC 7380, NGC 6871, and Hogg 10. The observed dependency of  $E(B - V)$  shows that early type stars in these clusters are having some amount of dust and gas around them.
5. From the near-IR flux values, we identified eight stars in Collinder 232, four stars in NGC 6530, and one star in NGC 6231 which are having significant near-IR excess fluxes. Higher fluxes may be due to the presence of circumstellar material around these stars.

## ACKNOWLEDGEMENTS

We are thankful to the anonymous referee for careful reading of the paper and constructive comments. We also thank Prof. Johan P. U. Fynbo, Niels Bohr Institute, University of Copenhagen, Denmark for reading our manuscript. This research has made use of the WEBDA database, operated at the Department of Theoretical

Physics and Astrophysics of the Masaryk University. We have also used data from Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

## REFERENCES

- Bhattacharjee, S. K., & Williams, I. P. 1980, MNRAS, 192, 841  
 Burki, G. 1975, A&A, 43, 37  
 Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, ApJ, 345, 245  
 FitzGerald, M. P. 1970, A&A, 4, 234  
 Fitzpatrick, E. L. 2004, in ASP Conf. Ser., Vol. 309, Astrophysics of Dust, eds. A. N. Witt, G. C. Clayton, & B. T. Draine, (San Francisco: ASP), 33  
 Johnson, H. L., & Morgan, W. W. 1953, ApJ, 117, 313  
 Johnson, H. L. 1966, ARA&A, 4, 193  
 Joshi, Y. C. 2005, MNRAS, 362, 1259  
 Hayashi, C. 1970, MSRSL, 19, 127  
 Kiszkurno, E., Kołos, R., Krelowski, J., & Strobel, A. 1984, A&A, 135, 337  
 Koornneef, J. 1983, A&A, 128, 84  
 Krelowski, J., & Strobel, A. 1983, A&A, 127, 271  
 Larson, B. 1973, ARA&A, 2, 219  
 Martin, P. G., & Whittet, D. C. B. 1990, ApJ, 357, 113  
 McCuskey, S. W., & Hauk, N. 1964, Asir. J., 69, 412.  
 McNamara, B. J. 1976, AJ, 81, 845  
 Nishiyama, S., et al. 2005, ApJ, 638, 839  
 Pandey, A. K., Mahra, H. S., & Sagar, R. 1990, AJ, 99, 617  
 Pandey, A.K., Upadhyay, K., Nakada, Y., & Ogura, K. 2002, A&A, 397, 191  
 Sagar, R. 1985, Abastumani astrophys. Obser. Mt. Kanobili Bull., 59, 191  
 Sagar, R. 1987, MNRAS, 228, 483  
 Sagar, R., & Qian, Z. Y. 1989, MNRAS, 240, 551  
 Sagar, R., & Qian, Z. Y. 1990, ApJ, 353, 174  
 Sagar, R., & Qian, Z. Y. 1993, BASI, 21, 565  
 Samson, W. B. 1975, Astrophys. Space 5d., 34, 363  
 Seab, C. G., & Shull, J. M. 1983, ApJ, 275, 652  
 Smith, R. G. 1987, MNRAS, 227, 943  
 Stone, R. C. 1977, A&A, 54, 803  
 Strom, K. M., Strom, S. E., & Yost, J. 1971, ApJ, 165, 479  
 Strom, K. M., Strom, S. E., Yost, J., Carrasco, L., & Grasdalen, G. 1972, ApJ, 173, 353  
 Tapia, M., Roth, M., Marrow, M., & Ruiz, M. T. 1988, MNRAS, 232, 661  
 Tapia, M., Costero, R., Echevarria, & Roth, M. 1991, MNRAS, 253, 649  
 Turner, D. G. 1994, RMxAA, 29, 163  
 Vashchinnikov, N. V., & Ilin, V. B. 1987, SvAL, 13, 157  
 Warner, J. W., Strom, S. E., & Strom, K. M. 1979, ApJ, 213, 427  
 Yadav, R. K. S., & Sagar, R. 2001, MNRAS, 328, 370  
 Yorke, H. W., & Krugel, E. 1977, A&A, 54, 183